

Abstract Submitted  
for the MAR08 Meeting of  
The American Physical Society

Sorting Category: 05.17.1 (E)

**Penetration depth study of Li, C and Li+C doped MgB<sub>2</sub> single crystals** CATALIN MARTIN, MATTHEW VAN-NETTE, RUSLAN PROZOROV, Ames Laboratory, Ames, IA 50011, J. KARPINSKI, N. ZHIGADLO, R. KHASANOV, Laboratory for Solid State Physics, ETH, 8093 Zurich, Switzerland — Magnetic penetration depth was studied in single crystals of MgB<sub>2</sub>, pure and doped with Li, C and Li+C. At zero applied field London penetration depth is measured and superfluid density,  $n_s$ , can be evaluated. We analyze effect of the dopings on  $n_s$ , thus on the superconducting gaps. In the vortex state, Campbell penetration depth is measured and it provides information about true critical current unaffected by the magnetic relaxation. In particular, penetration depth in vortex state becomes strongly irreversible, and we show that C doping enhances the irreversibility region, whereas the substitution with Li does not affect it. The results are discussed in terms of two-gap nature of MgB<sub>2</sub>.

☒ Prefer Oral Session  
☐ Prefer Poster Session

Catalin Martin  
cmartin@ameslab.gov  
Ames Laboratory

Date submitted: 19 Nov 2007

Electronic form version 1.4